

Lucien and Marguerite d’Azambuja, explorers of solar activity (1899-1959)

J.-M. Malherbe, Observatoire de Paris, PSL Research University, LESIA, Meudon, France

Email : Jean-Marie.Malherbe@obspm.fr

Date: 30 January 2024

ORCID identification : <https://orcid.org/0000-0002-4180-3729>

ABSTRACT

Henri Deslandres initiated imaging spectroscopy of the solar atmosphere in 1892 at Paris observatory. He developed two kinds of spectrographs: the “*spectrohéliographe des formes*”, i.e. the monochromatic instrument to reveal chromospheric structures (such as filaments, prominences, plages and active regions); and the “*spectrohéliographe des vitesses*”, i.e. the spectroheliograph to record line profiles of cross sections of the Sun, in order to measure the Dopplershifts of dynamic features. Deslandres moved to Meudon and hired Lucien d’Azambuja in 1899; they together built the large quadruple spectroheliograph. Call K systematic observations of the full Sun started in 1908 and were followed in 1909 by H α with two dedicated 3-metre spectroheliographs. Daily observations were organized by d’Azambuja who also intensively used the large 7-metre spectroheliograph for his research and thesis (1930). This paper summarizes fifty years of investigations by Mr and Mrs d’Azambuja, who explored various photospheric and chromospheric lines, performing special spectroheliograms with the high dispersion 7-metre instrument. They also observed intensively filaments and prominences, reported results in an extensive memoir (1948) and recorded rare solar activity events with the two 3-metre spectroheliographs, until their retirement in 1959.

KEYWORDS

d’Azambuja, solar physics, photosphere, chromosphere, filaments, prominences, spectroscopy, spectroheliograph, spectroheliogram, Dopplershifts

1 - INTRODUCTION

Henri Deslandres (1853-1948, see biography by Lequeux, 2023), defended his “*Doctorat ès Sciences*” (1888) in the laboratory of Alfred Cornu at Ecole Polytechnique, and was hired in 1889 by Admiral Ernest Mouchez (the director of Paris observatory since 1878). He was in charge of organizing a spectroscopic laboratory, in the context of the development of physical astronomy initiated by Janssen (1824-1907) at Meudon (this is the physics and chemistry of celestial objects). Deslandres first built a classical spectrograph to probe the solar atmosphere [note 1] using photographic means; he studied the line profiles of the ionized Calcium at 3934 Å wavelength (the Call K line) [note 2]; he succeeded to resolve the fine structure of the core in 1892. This was the starting point of the great adventure of spectroheliographs in France. With a classical spectrograph, monochromatic images can be delivered by an output slit located in the spectrum (selecting the light of a spectral line) when the input slit scans the solar surface. This is possible by moving the solar image upon the first slit (Janssen, 1869). The photographic spectroheliograph was invented on this basis by George Hale (1868-1938) in Kenwood (1892) and by Henri Deslandres in Paris (1893), simultaneously but independently. The full story was described in details by Malherbe (2023).

Many spectroheliographs were built in the world, following Hale and Deslandres techniques. For instance, long series of continuous observations in the Call K line were collected by the spectroheliographs of Kodaikanal in India, Mount Wilson in the USA, Mitaka in Japan, Sacramento Peak in the USA, Coimbra in Portugal, Meudon in France or Arcetri in Italy. They produced extended archives, some of them covering up to 10 eleven-years solar cycles, which are convenient to investigate long-term solar activity and study rare events (such as energetic flares, huge sunspot groups or giant filaments [note 3]). The Call K3 and H α central intensities are very convenient for imaging spectroscopy of the chromosphere (filaments, prominences, magnetized bright plages, altitude range 1500-2000 km); the wings of the Call K line (K1v) formed in the lower range (200-500 km) are good tracers of sunspots and faculae [note 3]. The Call K line is mainly sensitive to temperature fluctuations, while H α is more sensitive to the density. In parallel, between 1919 and 1939, “section” spectroheliograms of Call K were systematically performed in Meudon. In that case, instead of moving continuously, the entrance slit of the spectrograph scanned the sun by steps of 20”-25”, forming on the photographic plate full spectra of cross sections of the Sun. The spectral width was about 2 Å, allowing to evaluate the Dopplershifts [note 4] of the Call K3 component (the line core). Such observations were unique, but the derivation of radial velocities was, of course, a purely manual and long task. For that reason, this

program was abandoned in 1939 at the entry of WW2, and never restarted later. However, the 2017 version of Meudon spectroheliograph, with a fast sCMOS numerical detector, registers the full line profiles and provides now 3D FITS data-cubes (x , y , λ), in the astronomical standard format, which are available on-line and allow Dopplershifts determination and more parameters using line inversion techniques.

This paper reviews briefly fifty years of monochromatic observations performed in Meudon observatory, during the first half of the twentieth century, under the auspices of Lucien d’Azambuja (1884-1970, [figure 1](#)), who retired in 1954. Section 2 describes systematic observations started in 1908. Section 3 relates the studies performed by Lucien and Marguerite d’Azambuja (born Roumens, 1898-1985, [figure 1](#)), forming together the famous team of filament and prominence explorers, until 1959. Section 4 summarizes special observations of various spectral lines, done before 1930 for d’Azambuja’s thesis with the 7-metre spectroheliograph, which demonstrated that the initial choice of Call K and H α was the best for imaging spectroscopy of the chromosphere. At last, section 5 presents rare events of solar activity observed with the more classical 3-metre spectroheliographs.



November 1939

Astronomers

- (1) M. d'Azambuja**
(42 years old)
- (2) L. d'Azambuja**
(55 years old)
- (3) B. Lyot**
- (4) R. Servajean**

Visiting astronomer

- (5) P. Phu**

- (6) Mrs Lyot**

- (7) Mrs Servajean**

Figure 1. Marguerite d’Azambuja, Lucien D’Azambuja and others in November 1939, the day of P. Phu thesis (courtesy Paris Observatory).

2 – SYSTEMATIC OBSERVATIONS AT MEUDON

Systematic observations started in 1908 for Call K and 1909 for H α with the new spectroheliographs installed in Meudon by Deslandres and d’Azambuja (Deslandres, 1910; d’Azambuja, 1920a, 1920b, 1930) as shown by [figures 2, 3, 4](#). Details of the optical setup were summarized by Lequeux (2022) and Malherbe (2023). A spectrohelioscope ([figure 5](#)) working in H α came in the fifties for visual inspection of solar activity. It was a removable device composed of a rotating prism in front of the entrance slit (to scan the Sun) and a second one at the output slit (to form a monochromatic image using retinal persistence with an eyepiece).




Figure 2. The two-mirror coelostat with Lucien d'Azambuja in 1921 (Courtesy Gallica/BNF, Meurisse agency).




Figure 3. The spectroheliographs with Lucien d'Azambuja in 1921 (Courtesy Gallica/BNF, Meurisse agency).




Figure 4. The spectroheliographs designed by Deslandres and d’Azambuja (1908). The entrance objective ($f = 4.0$ m, **a**) delivers a 37.2 mm image and scans the Sun with the carrier (**C**), velocity transformer (**T**), motor (**M**). The spectrograph slit (**b**), the collimator ($f = 1.3$ m, **c**) are common to all configurations. Top: spectroheliographs for systematic observations in Call K (three prisms **g**, left panel) and H α (grating **d**, right panel). Both use a 3 m focal length chamber with an output slit in the spectrum (**i**, **f**). The photographic glass plate holders are motorized and move 2.31 times faster (the optical magnification) than the imaging objective. The diameter of the Sun is 86 mm on plates. Bottom: the high dispersion spectroheliograph for research. It uses either the grating (**d**) or prisms (**g**). **h** is a flat mirror. **l** is a 7 m concave chamber that forms a spectrum in (**m**). The diameter of the Sun is there 205 mm. After comes an afocal reducer composed of a second 7 m concave mirror (**n**) and lenses (**p**, **q**, **r**) providing various magnifications on the photographic plate and output slit (**s**). There is a prism (**o**, 30°) to reject parasitic light or unwanted orders.




Figure 5. The visual spectrohelioscope in the fifties with L. d’Azambuja in the stamp (courtesy Paris Observatory).




Figure 6. Formation heights (in abscissa) of various solar spectral lines. The cores of H α and Ca II K (named K3) form respectively at about 1500 km and 1900 km in the chromosphere (the layer from 500 km to 2100 km with increasing temperature). The Ca II K2 peaks ($K3 \pm 0.2 \text{ \AA}$) form in the low chromosphere (1000 km), while the wings of H α and Ca II K (named K1, i.e. $K3 \pm 1.3 \text{ \AA}$) form respectively at about 200 km and 500 km in the photosphere (the 500 km layer above the surface with decreasing temperature). The Mg II k line in UV (observed by the IRIS/NASA spectrograph in space) looks like Ca II K (after Vernazza et al, 1981).




Figure 7. Typical spectroheliograms of the period 1919-1939 (example of 11 July 1937). Call K1v for the photosphere (sunspots, faculae); Call K3 for the chromosphere (filaments, bright plages); Call K line profiles (cross sections of the Sun by 22'' steps for Dopplershift measurements); H α for the chromosphere (filaments, plages). Courtesy Paris observatory.

Systematic observations were performed with two 3-metre spectroheliographs (figures 3, 4 and movie 1): spectrograph n°I for Call K using 3 prisms of 60° at minimum deviation, providing a spectral resolution of 0.15 Å, and spectrograph n°II for H α using a Rowland plane grating, 568 grooves/mm, providing in the first order a spectral resolution of about 0.40 Å. The focal length of the scanning objective was 4.0 m, and the magnification of the spectrographs was 2.31 (3.0 m chamber, 1.30 m collimator), so that the size of the Sun on the photographic glass plates was 86 mm. The width of the entrance slit was 0.03-0.04 mm (1.5"-2") well adapted to the usual seeing of 2" in Meudon; while the second slit in the spectrum was 0.075 mm wide.

From 1908 to WW1 (1914), observations consisted of daily spectroheliograms in Call K3 (line centre, chromosphere), Call K1v (violet line wing, photosphere) and H α centre (chromosphere). Formation heights of K3, K1 and H α lines are reported in figure 6. Observations were totally interrupted during WW1. Figure 7 shows typical observations performed daily from 1919 to 1939 until the start of WW2. During this period, Call K3, Call K1v and H α were continued, and a "section" spectroheliogram (figure 8) with an enlarged slit in the spectrum (1 mm = 2 Å) was added for Dopplershift measurements of the Call K line; for that purpose, full spectra of 86 cross sections of the solar surface were recorded by spatial steps of about 1 mm = 22". Contrarily to WW1, observations were not stopped by WW2, but the lack of manpower was an important constraint, so that "section" spectroheliograms were abandoned, and never restarted after WW2. Typical systematic observations done after WW2 by the d'Azambujas (figure 9) are shown in figure 10; they still include the initial choice of 1909 (Call K3, K1v and H α), plus a long exposure Call K3 for limb features and prominences, with a disk attenuator (because prominences are 5-10 times fainter than the disk).




Figure 8. Typical example of a "section" spectroheliogram in Call K line, 17 September 1938. The imaging objective was moving by steps of about 22". The slit in the spectrum was enlarged from 0.075 mm to 1.0 mm, in order to pass 2.0 Å instead of 0.15 Å, the usual bandwidth for Call K3 and Call K1v images of the chromosphere. Each of the 86 cross sections provides the line profiles; the Dopplershifts of the line core (K3) were measured and results reported as an index (0-5) in the tables accompanying the synoptic charts of the chromospheric structures, started in 1919 by d'Azambuja for each rotation. Since the Dopplershift determination was very time consuming for observers, which were not numerous, this work was done during twenty years (1919-1939) and was abandoned before WW2. Courtesy Paris observatory.

Systematic observations continued under the direction of Lucien d'Azambuja until his retirement in 1954 (he was 70 years old, Meudon astronomers were those of [figure 9](#) at that epoch). At this date, Mrs d'Azambuja became in charge of observations, but his husband was still working as a volunteer; she retired in 1959 (after 60 years of astronomy for L. d'Azambuja !). Gualtiero Olivieri (1928-2018) and Marie-Josèphe Martres (1924-2014) became respectively responsible of observations and of the synoptic map program initiated in 1919 by L. d'Azambuja (synthetic drawings of solar activity for each solar rotation, showing the average position of filaments, sunspots and facular regions).



Astronomers in 1950:

- (1) L. d'Azambuja
- (2) M. d'Azambuja
- (3) H. Grenat
- (4) A. Dollfus
- (5) F. Baldet
- (6) R. Herman

Figure 9. Scientific people working at Meudon Observatory in 1950 (Courtesy Paris Observatory).

All data since 1908 are freely available on-line at <https://bass2000.obspm.fr>, in 12 bits FITS and 8 bits JPEG format, for images after 1980, and in JPEG only for older spectroheliograms (250 dpi scans of glass and film plates). A new scan at 600 dpi/16 bits is in progress and should be completed in 2025. The collection is one of the longest world-wide, with more than 100000 images along 10 solar cycles.

The spectroheliograph dedicated to daily observations was revisited in the eighties. The prisms (for Call K) and plane grating at order 1 (for H α) were replaced by a unique 300 groves/mm grating (blaze angle of 17°27') providing alternatively H α at order 3 and Call K at order 5, without any rotation, so that the same 3-metre chamber was used for both lines. The order 4 was convenient for H β but the lenses were achromatized only for orders 3 (red) and 5 (violet). Interference orders were selected by coloured filters. Glass plates were abandoned and systematic observations were produced in this new configuration with 13 x 18 cm² film plates until year 2000. Almost real time digitization of films started in 1995 with a 12 bits scanner and images were archived by the BASS2000 data base when it was commissioned in 1996.

The second slit in the spectrum was removed in 2001 when film plates were replaced by a back illuminated CCD camera (1340 x 100 format) from Princeton Instruments (14 bits dynamic range, 20 μ m pixels), reducing the chamber focal length from 3.0 m to 0.9 m. Finally, full line profiles are recorded since 2017 with a new scientific CMOS camera from PCO (16 bits dynamic range, 2048 x 2048 format, 6.5 μ m pixels) for each location (x, y) of the solar surface, with a chamber reduced to 0.40 m only. As Call H and Call K are simultaneously observed with this sCMOS array, both lines are recorded ([movie 2](#) for x-direction surface scans and wavelength exploration of final Call K and H α data-cubes). The spectral sampling of Call H and K is 0.093 Å (order 5), while the one of H α is 0.155 Å (order 3). The spectral domain of usual observations is 9 Å and 6 Å, respectively for Call H, K, and H α . The spatial sampling is about 1.1", which is quite convenient for the standard Meudon seeing of 2". The width of the entrance slit of the spectrograph never changed (0.03 mm) and corresponds to 1.55" on the solar surface.




Figure 10. Typical spectroheliograms after WW2 (example of 28 October 2003, the day of a massive X-class flare). *H α* for the chromosphere (filaments, plages); *CaII K1v* for the photosphere (sunspots, faculae); *CaII K3* for the chromosphere (filaments, bright plages); *CaII K3* long exposure time for prominences and limb with a neutral density attenuator ND1 (10% transmission) superimposed on the solar disk. Courtesy Paris observatory.

3 - Mr AND Mrs d'AZAMBUJA, THE FAMOUS PROMINENCE EXPLORERS

Lucien d'Azambuja was hired in 1899 by Deslandres when he was only 15 years old (Rösch, 1970, Martres, 1998). Of course, he completed in parallel his studies in mathematics and physics at Paris University. Marguerite Roumens arrived at Meudon in 1925 and participated to the spectral line observing program of d'Azambuja (section 4) with the 7-metre spectroheliograph; the original results were reported in his thesis (d'Azambuja, 1930). They got married and started a common and monumental research work about solar filaments (d'Azambuja & d'Azambuja, 1948). This memoir is considered as a “bible” among Meudon solar astronomers. It contains an extensive study of the morphology, evolution and stability of solar filaments and prominences (figure 11), and their relationships with their neighbourhood (active regions). It is a founding work at the base of modern solar physics. When L. d'Azambuja retired in 1954, his wife became responsible of observations; she retired in 1959 and the d'Azambujas left together Meudon at this date, after 60 years of astronomy for Mr d'Azambuja (Malherbe, 2023) ! An incredible career, impossible to reproduce today...




Figure 11. Various aspects of solar prominences in $H\alpha$ and Call K, observed with Meudon spectroheliographs during the first half of the twentieth century (false colours, rainbow coding). Courtesy Paris observatory.




Figure 12. A solar filament exhibiting various perspective effects during the rotation (27 August and 3 September 1929), in Call K3 (left) and $H\alpha$ (right). Courtesy Paris Observatory.

This huge study of 387 filaments showed that they are thin (5000 km typical), long (10^5 km to 10^6 km), high (up to 50000 km) and cold (8000 K) Hydrogen structures suspended in the hot corona (10^6 K), above the chromosphere. Filaments look like a vertical curtain anchored into the photosphere by legs and barbs on both sides. They exhibit extremely variable aspects due to perspective effects during the solar rotation (figure 12 and figure 13). When seen at the limb, filaments are called bright prominences which emit light in $H\alpha$ or Call K. But when seen on the solar disk, filaments absorb the photospheric light in the same lines and appear dark.




Figure 13. Various aspects of solar filaments during the rotation. Emitting light at the limb (bright prominences), or absorbing light on the disk (dark filaments). After d’Azambuja & d’Azambuja (1948).




Figure 14. Evolution of filaments and their anchorage into the photosphere (arrows indicate barbs along the filament and legs at both ends). After d’Azambuja & d’Azambuja (1948).

Figure 14 shows a typical example of filament evolution during a few days. D’Azambuja & d’Azambuja (1948) found that quiescent filaments (far from active regions) evolve slowly and can last during several rotations (the duration of the synodic or Carrington rotation is 27.27 days, it corresponds to the rotation of zones at $\pm 26^\circ$ latitude, seen from the Earth). Their shape and length vary continuously, filaments can split into

several parts, which can also reconnect. Varying perspective effects play a major role during the solar rotation. The authors found a systematic inclination (8°) of the filament blade towards the western direction. The orientation of equatorial filaments is not far from the meridians (figure 14), while polar ones, which appear at the beginning of a new cycle, are close to the parallels (figure 15). The differential rotation (faster at low latitudes) affects the direction of long filaments and elongate them. We know today that filaments are dense (density of 10^{17} m^{-3}) and cold structures (8000 K) supported in the tenuous (10^{15} m^{-3}) and hot corona (10^6 K) by weak magnetic fields (0.002 T) overlying the boundary between regions of opposite polarities. At the beginning of a new cycle, polar filaments delineate the frontier between the new emerging magnetic field at lower latitudes and the old polar one which reverses at the next solar maximum. The authors emphasized the importance of royal zones, both for the birth of active regions and filaments; they discovered that filaments migrate towards the poles, indicating the existence of a meridional circulation, which was confirmed later by helioseismology (the MDI instrument onboard SOHO/ESA/NASA).




Figure 15. Evolution during the solar rotation of a polar prominence (February 1938). Father Angelo Secchi (1818-1878) wrote in his book *“Le Soleil”*: “Prominences exhibit so various shapes that it is impossible to describe them with any exactitude”... Courtesy Paris Observatory.

The rotation of filaments was found similar to the one of active regions and sunspots, except at high latitudes where measurements revealed a slower rotation. Among the 387 studied filaments, 141 were associated to active regions (plage filaments), while 246 were not (quiescent ones). Their temporal evolution was found similar: they need (in average) three rotations to develop, and start to fragment during the fourth passage on the disk. They can last up to six rotations (even twenty rotations for very quiet polar prominences) and their length increases with their lifetime. Some filaments were observed to split or merge together. They may become unstable. During a full solar cycle (11 years), D’Azambuja & d’Azambuja (1948) counted that 206 objects partly disappeared (this was called the “Disparition Brusque” phenomenon, or DB); while in 137 cases, the DB was only temporary. Most perturbations occurred among equatorial filaments close to active regions. We know today that the DB events can be either thermal (heating of material which vanishes in $H\alpha$) or dynamic (magnetic instability affecting the filament support, sometimes associated to flares or coronal mass ejections).

4 - SPECIAL SPECTROHELIOGRAMS AND UNUSUAL SPECTRAL LINES

During his thesis work, that he defended in 1930, Lucien d’Azambuja explored many spectral lines with the large 7-metre spectroheliograph for research, providing a much higher dispersion than the two 3-metre spectroheliographs used for systematic observations (figure 16). High dispersion is required to study thin photospheric lines such as $\text{MgI } 3838 \text{ \AA}$, $\text{SrII } 4078 \text{ \AA}$, $\text{FeI } 4202 \text{ \AA}$, $\text{CaI } 4227 \text{ \AA}$, $\text{FeI } 4384 \text{ \AA}$, $\text{MgI } 5184 \text{ \AA}$, $\text{HeI } 5876 \text{ \AA}$, $\text{NaI } 5890 \text{ \AA}$, as well as chromospheric infrared lines of $\text{CaII } 8498 \text{ \AA}$, $\text{CaII } 8542 \text{ \AA}$, $\text{HeI } 10830 \text{ \AA}$. The large spectroheliograph (dismantled in the sixties) was able to offer the dispersion of 1 mm/\AA , allowing excellent spectral resolutions on line profiles of about 0.05 \AA . Prisms were used in the blue part of the spectrum, and a plane grating for longer wavelengths.




Figure 16. Dispersion of the various spectroheliographs. 3-metre spectrographs n°I (prisms) and n°II (grating) were used only for systematic observations, respectively for CaII K and H α lines. The high dispersion 7-metre spectroheliograph n°IV (3 prisms or plane grating) was used only for research work. After d'Azambuja (1930).




Figure 17. Spectral lines of CaII K and H α above a bright plage (top), at the limb (top) or in the quiet Sun (bottom). Wavelength in abscissa. After d'Azambuja (1930).

For instance, the high dispersion 7-metre spectroheliograph provided the results of [figure 17](#) for CaII K and H α lines, with a lot of details along the profiles. In plages or at the limb, the bright CaII K2v and K2r

peaks ($\pm 0.2 \text{ \AA}$ apart from the dark K3 core) are well resolved. The chromosphere appears clearly at the limb in both lines above the photosphere (which is visible in the line wings and in the continuum).

Figure 18 displays three images, respectively in the centre of $H\alpha$ line and in the blue wing. Filaments are still visible at -0.25 \AA , but vanish at -0.50 \AA (near the inflexion points of the line profile) and the photosphere (below the chromosphere) begins to appear at -0.50 \AA , except in the case of Dopplershifts ($0.5 \text{ \AA} = 25 \text{ km/s}$). Such observations are interesting because they were never done daily, contrarily to the $H\alpha$ centre.




Figure 18. Observations at three positions in the $H\alpha$ line. Filaments vanish in the line wing, except in the case of Dopplershifts, while photospheric structures begin to appear. 5 June 1930. After d’Azambuja (1930).




Figure 19. Observations at three positions in the Call K line (K3, K2v, K1v). 29 July 1927. After d’Azambuja (1930).

Figure 19 displays three images, respectively in the centre of Call K line (K3) and in the blue wing. Filaments and bright plages are perfectly visible in K3, but this is not the case of sunspots. Filaments are fairly observable at -0.25 \AA (K2v) and sunspots begin to appear. But filaments completely vanish at -1.27 \AA (K1v) while the photosphere, below the chromosphere, becomes clearly visible, with sunspots and bright faculae (which are located just below the chromospheric plages). Observations of K2v are particularly rare in the collection, contrarily to K3 and K1v.

Figure 20 shows an exceptional observation of the Balmer series, $H\alpha$, $H\beta$, $H\gamma$, $H\delta$ and $H\epsilon$. The contrast of filaments and plages (chromosphere) vanishes with decreasing wavelength, because lines form deeper. On the contrary, sunspots appear progressively along the series, which reveals that $H\alpha$ is the best choice for chromospheric structures, while $H\delta$ or $H\epsilon$ are not better than Call K1v for sunspots and faculae (however $H\epsilon$ is contaminated by the red wing H1r of the Call H line). This observational series is the only one of the collection. Figure 21 focuses on the central part of the solar disk with active regions and filaments.




Figure 20. The Balmer series $H\alpha$, $H\beta$, $H\gamma$, $H\delta$ and $H\epsilon$ of Hydrogen, 14 August 1947, and the Call K3 spectroheliogram for comparison. $H\alpha$ is chromospheric, but $H\delta$ and $H\epsilon$ are rather photospheric. Courtesy Paris Observatory.




Figure 21. The Balmer series $H\alpha$, $H\beta$, $H\delta$ and $H\epsilon$, 14 August 1947 (detail of [figure 20](#)). The contrast of filaments and plages is the best in $H\alpha$ and decreases drastically along the series. After d'Azambuja & d'Azambuja (1948).

Helium lines were also tested by d'Azambuja. Figure 22 shows an exceptional observation in Hel D3 5876 Å of an active region producing the huge flare of 26 July 1946, while figure 23 shows the first world-wide observation done in the infrared Hel 10830 Å line. These observations are absolutely unique in the collection (d'Azambuja, 1938). Several decades later, Kitt Peak National Observatory started systematic observations in this infrared line of Helium.




Figure 22. The flare of 25 July 1946 observed in Hel D3 at 5876 Å from 16:58 to 17:03. Courtesy Paris observatory.




Figure 23. The first observation of the infrared Hel 10830 Å line (right), where both filaments and plages appear darker than the quiet Sun. Call K3 (left) is given for comparison. 17 September 1938. Courtesy Paris observatory.

Infrared lines of Call were studied by d’Azambuja in his thesis work. [Figure 24](#) shows images in Call 8498 Å and 8542 Å, on 4 September 1928, in comparison with the usual Call K3 3934 Å. These observations are extremely rare in the collection and showed that chromospheric structures are better seen with Call K3 ([figure 25](#)). Several decades later, Kitt Peak National Observatory started systematic observations in Call 8542 Å in order to measure chromospheric magnetic fields (which is much more difficult with Call K).

[Figure 25](#) shows also a detail of [figure 23](#), for comparison of Hel 10830 Å and Call K. While filaments appear dark in both lines, this is not the case of plages, which are bright in Call K but dark in Hel 10830 Å.




Figure 24. Call 8498 Å and Call 8542 Å, compared to Call K3 3934 Å (right). 4 September 1928. After d’Azambuja (1930).




Figure 25. Call 8542 Å, compared to Call K3 3934 Å (top). 29 June 1928. After d’Azambuja (1930). Hel 10830 Å, compared to Call K3 3934 Å (bottom). 17 September 1938. Courtesy Paris Observatory.

Cal 4227 Å line was studied by d’Azambuja during the course of his thesis. Figure 26 shows images in the line core and wing, in comparison with the usual Call K1v for the photosphere. Figure 27 shows spectroheliograms in FeI 4046 Å, FeI 4202 Å, SrII 4078 Å lines. Many more photospheric lines (the list of figure 27 indicating also the selected bandwidth in Å) were observed at high spectral resolution and a facular visibility index (0-5) was established. It sounded clearly that these investigated lines provided spectroheliograms that are comparable to Call K1v, so that observations in these wavelengths were never reproduced later.




Figure 26. Spectroheliograms of the Cal 4227 Å line, compared to Call K1v for the photosphere (at right, for reference, the chromospheric Hα line). After d’Azambuja (1930).




Figure 27. Spectroheliograms in various photospheric lines, compared to Call K1v. After d’Azambuja (1930).

5 - RARE EVENTS OF SOLAR ACTIVITY OBSERVED WITH THE SPECTROHELIOGRAPHS

The centennial Meudon collection of Call K1v spectroheliograms contains plenty of information about the area covered by sunspots. In particular, the April 1947 sunspot group (figure 28, centre) is probably the largest ever seen (6.1% of the solar disk), followed by other groups of the same solar cycle (number 18), among them the February 1946, May 1951, July 1946 (figure 28) and March 1947 groups (respectively 5.2%, 4.9%, 4.7% and 4.6% of the disk). These observations illustrate the interest for long term observations spanning more than 10 solar cycles, because exceptional area sunspots provide an upper limit for magnetic energy that can be released in solar flares. Please refer to Malherbe (2023) for more details on such events.




Figure 28. Huge sunspot groups (25 July 1946, 5 April 1947, 17 May 1951 seen in Call K1v). Courtesy Paris observatory.

Extremely long filaments, such as those of figure 29 and figure 30, delineate giant magnetic cells on the Sun. Indeed, filaments are located above the inversion line of photospheric magnetic fields; large filaments can be considered as tracers of large magnetic zones. Filaments of more than one solar radius length are not frequent and can last more than 6 rotations (figure 30). Here again, long term collections of observations, are essential to catch rare phenomena which contribute to a better understanding of solar physics.




Figure 29. Long filaments of 26 May 1930 in Call K3 (left) and H α (right). After d’Azambuja & d’Azambuja (1948).

Fast solar activity events, such as flares, “Disparitions Brusques” of filaments, prominence eruptions and mass ejections are difficult to observe with a spectroheliograph, because of the moderate temporal resolution. For that reason, H α filters were used after 1950. At Meudon, high cadence observations (1 minute)

of dynamic events started in 1954 with Lyot filters, in particular in the frame of the International Geophysical Year (1957). However, until this date, spectroheliographs were the only available instruments to study fast evolving phenomena, and a few exceptional active phenomena were recorded with a maximum of twelve spectroheliograms for an event lasting a few hours. It was indeed a complicated task at the epoch of glass photographic plates. The prominence eruption of 18 June 1925 ([figure 31](#) and [figure 32](#)) is one of the best events available in the collection. A “section” spectroheliogram ([figure 32](#)) was even produced to estimate the radial velocities (i.e. the velocity component orthogonal to the sky plane). Of course, prominence eruptions are today commonly observed in HeII 304 Å (the Ly α line of HeII at 80000 K) with the AIA telescope onboard SDO/NASA since 2010 (45 s cadence for each waveband), but it is also the case with the EIT telescope onboard SOHO/ESA/NASA since 1996 (with a lower cadence and spatial resolution than SDO).




Figure 30. An extremely long filament observed during several rotations. It started to form during rotation 915 (not shown). The length at rotation 920 exceeds one million kilometres. After d’Azambuja & d’Azambuja (1948).




Figure 31. Instability of a solar prominence at the limb observed in Call K3 on 18 June 1925 (08:46 to 11:23 UT) with Meudon spectroheliograph; an artificial moon was masking the solar disk. Courtesy Paris observatory.




Figure 32. Instability of the solar prominence of 18 June 1925 in the Call K line from 09:11 to 11:03 UT. The image at left is a “section” spectroheliogram, obtained by moving the entrance slit by steps of 22”. The five line profiles of Call K (2.15 Å wide) at the bottom are those of the cross section AB. Courtesy Paris observatory.

Figure 33 displays one of the “treasures” of the 10 solar cycles Meudon collection, this is the eruption of a huge solar prominence observed just after WW1 in May 1919 (after a total interruption during the war). Observations were performed in the Call K3 line with an artificial moon superimposed upon the solar disk to allow long exposure time, as prominence material is faint.

The July 1946 group of figure 28 produced the strong solar flare of figure 34, and a huge geomagnetic storm. This phenomenon is one of the most energetic eruptions ever observed. André Danjon, the director of Paris-Meudon observatories, was there during this event (he often spent the summer in Meudon) and was fond of solar observations. However, Gualtiero Olivieri, who was observer, reported that “his advices and commands given to the observers did not helped them to stay quiet and keep cool” ! Flares originate in active regions and have a magnetic origin. The magnetic energy is stored in sunspot groups, such as the big ones of figure 28; it is proportional to the surface and to the square of the magnetic field intensity. Unstable configurations often occur near the solar maximum and just after, and trigger huge flares, which convert magnetic to kinetic energy and heat. The 1946 event illustrates the importance of historical data. It allowed to predict the maximum energy of a solar flare, about $5 \cdot 10^{26}$ Joule (10 times the one ever measured by satellites of the modern era, Aulanier *et al*, 2013). The average energy of solar eruptions lies around 10^{25} Joule.




Figure 33. Instability of a solar prominence seen in Call K3 on 27, 28 and 29 May 1919, observed with Meudon spectroheliograph. The times are indicated in the picture. Courtesy Paris observatory.




Figure 34. The major two-ribbon solar flare of 25 July 1946 observed in H α with Meudon spectroheliograph (the times are, from the right to the left, 15:37, 16:14, 16:24, 17:57, 18:15 UT). The leftmost image at 16:58 UT was got in the HeI D3 5876 Å line. The short-duration flash phase (lasting minutes) occurred at 16:14 UT. Two bright ribbons (17:57, 18:15 UT) formed during the long-duration gradual phase (lasting hours) and corresponded to the impact of energetic particles on the chromosphere. Courtesy Paris observatory.

6 – CONCLUSION

Meudon spectroheliographs produce monochromatic images of the chromosphere in Call K3 and H α , and of the photosphere in the blue wing (K1v) of the Call K line, on a daily basis since 1908. The collection extends over 10 solar cycles and contains more than 100000 spectroheliograms freely available at <https://bass2000.obspm.fr>. However, between 1919 and 1954, a lot of special and unique observations were produced by Mr and Mrs d’Azambuja with the large 7-metres spectroheliograph dedicated to research. Many unusual spectral lines were studied, such as photospheric lines of FeI, CaI, HeI or SrII, and chromospheric lines of infrared Call or HeI. Some observations, such as HeI 10830 Å, were the first world-wide, and showed that the initial choice of Call K3, K1v and H α was the best for imaging spectroscopy of the photosphere and the chromosphere. Besides, many dynamic and rare events were recorded with the spectroheliographs dedicated to systematic observations, such as huge flares or prominence ejections. Of course, spectroheliographs were not designed for high cadence phenomena, as their goal was low cadence and long term observations of the solar cycles. For that reason, instruments using Lyot H α filters were used at Meudon after 1954 to record fast evolving events.

7 - ACKNOWLEDGEMENTS

The author is indebted to I. Bualé and F. Cornu for spectroheliograms of the Meudon solar collection, and for the photographic archive of the instruments and characters.

8 - NOTES

[1] The solar atmosphere. It is composed of three layers: (1) the photosphere (the visible surface, temperature decreasing from 6000 K to 4500 K in 300 km) with dark magnetized sunspots and bright faculae around, (2) the chromosphere above (temperature increasing from 4500 K to 8000 K in 2000 km) with dark filaments and bright plages (corresponding to faculae in the photosphere), (3) the ionized and hot corona (2 million K). The corona extends at million kilometres and gives rise to the solar wind (charged particles), which propagates into the interplanetary medium. The chromosphere and the corona require spectroscopic means to reveal their structures, respectively via absorption lines of the visible spectrum or emission lines of the ultraviolet spectrum. The solar atmosphere follows a 11-year activity cycle and a 22-year magnetic cycle. Flares and coronal mass ejections occur in active regions a few years around the solar maximum; the next maximum of the current cycle (number 25) is forecasted for 2025. The symbol K above (Kelvin) is the unit of the absolute temperature, related to the Celsius temperature by $T(K) = T(^{\circ}C) + 273.15$.

[2] The wavelength. Electromagnetic waves, such as the visible light, are detected by the telescopes. They form a continuous spectrum (such as the rainbow) with superimposed spectral lines, revealing the presence of atoms, ions or molecules. In the visible or ultraviolet spectrum, the lines are identified by their wavelength, measured in nanometres ($1 \text{ nm} = 10^{-9} \text{ m}$), but spectroscopists often prefer the Angström ($1 \text{ Å} = 0.1 \text{ nm} = 10^{-10} \text{ m}$). The visible solar spectrum (4000 Å - 7000 Å) reveals thousands of absorption lines, while emission lines occur in the ultraviolet spectrum (below 4000 Å) or at the limb.

[3] The solar structures. Dark sunspots are regions of intense magnetic fields (0.1 - 0.3 T). Bright faculae or plages form, together with sunspots, active regions, and exhibit smaller magnetic fields (0.01 - 0.05 T). Dark filaments, also called prominences when seen at the limb, are thin and high structures (50000 km) of dense material suspended in the corona by weak magnetic fields (0.001 T). The symbol T is the unit of magnetic field (Tesla); the Gauss ($1 \text{ G} = 10^{-4} \text{ T}$) is also used. Flares occur in zones of unstable fields; reconnections convert magnetic energy into kinetic energy (ejections), radiation (X-rays) and heat (brightenings).

[4] The Doppler effect. When a light source moves in the observer’s direction, the spectral lines are shifted towards shorter wavelengths (blueshift). On the contrary, if the source moves in the opposite direction, lines are shifted towards longer wavelengths (redshift). The Dopplershift $w(\text{Å})$ is a wavelength shift, proportional to the projection V (km/s) of the velocity vector along the line-of-sight (it is also called radial velocity, positive towards the observer in solar physics): $w = -\lambda (V / C)$, where λ is the line wavelength (Å) and C is the speed of light ($3 \cdot 10^5 \text{ km/s}$).

9 – MPEG4 MOVIES

Movie 1: the Meudon spectroheliograph in 1942 (courtesy Paris Observatory)

Movie 2: scanning the Sun today with the numerical spectroheliograph (courtesy Paris Observatory)

10 - REFERENCES

- G. Aulanier, P. Démoulin, C.J. Schrijver, M. Janvier, E. Pariat and B. Schmieder, "The standard flare model in three dimensions – II – Upper limit on solar flare energy", 2013, *Astrophysical Journal*, 549, A66.
- L. d’Azambuja, "Le spectrohéliographe et les photographies quotidiennes de l’atmosphère du soleil. Part 1", 1920a, *L’Astronomie*, 34, 433-446.
- L. d’Azambuja, "Le spectrohéliographe et les photographies quotidiennes de l’atmosphère du soleil. Part 2", 1920b, *L’Astronomie*, 34, 487-498.
- L. d’Azambuja, "Recherches sur la structure de la chromosphère solaire", thesis, 1930, *Annales de l’Observatoire de Paris, section d’astrophysique, à Meudon*, volume 8, fascicule 2, Gauthier-Villars editor, Paris. On-line at Gallica/BNF. <https://gallica.bnf.fr/ark:/12148/bpt6k6537953n>
- L. d’Azambuja and M. d’Azambuja, "Images Nouvelles de la chromosphère solaire obtenues au spectrohéliographe avec la raie infrarouge de l’hélium 10830 et avec la raie H ϵ de l’hydrogène", 1938, *Bulletin Astronomique*, 11, 349-359.
- L. d’Azambuja and M.d’Azambuja, "Etude d’ensemble des protubérances solaires et de leur évolution, effectuée à l’aide des spectrohéliogrammes de Meudon et des cartes synoptiques publiées par l’établissement sous l’égide de l’UAI", 1948, *Annales de l’Observatoire de Paris, section d’astrophysique, à Meudon*, volume 6, fascicule 7, Gauthier-Villars editor, Paris. On-line at Gallica/BNF. <https://gallica.bnf.fr/ark:/12148/bpt6k65382116>
- H. Deslandres, "Recherches sur l’atmosphère solaire. Photographie des couches gazeuses supérieures. Etude simultanée des formes et des mouvements des vapeurs". 1910, *Annales de l’Observatoire d’Astronomie Physique de Paris, sis parc de Meudon*, tome 4, Gauthier-Villars editor, Paris. On-line at Gallica/BNF. <https://gallica.bnf.fr/ark:/12148/bpt6k65373577>
- J. Janssen, "Méthode pour obtenir des images monochromatiques des corps lumineux", 1869, *British Association for the Advancement of Science*, 39, 23.
- Lequeux, J., "Henri Deslandres: from molecular to solar physics", 2022, *JAHH*, vol 25, issue 3, 553-575.
- Lequeux, J., "Life at Meudon Observatory, 1876-1963", 2022, *JAHH*, vol 25, issue 2, 171-186.
- J.-M. Malherbe, I. Bualé, F. Cornu, D. Crussaire and T. Corbard, "The potential of Meudon spectroheliograph for investigating long term solar activity and variability", 2023, *Advances in Space Research*, 71, 1922, on-line at <https://doi.org/10.1016/j.asr.2022.07.058>
- J.-M. Malherbe, "130 years of spectroheliograms at Paris-Meudon observatories (1893-2023)", 2023, *JHA*, volume 54, issue 3, 274, also on-line at <https://arxiv.org/abs/2301.11105>
- M.-J. Martres, "Lucien d’Azambuja", 1998, *IAU Symposium 167, Astronomical Society of the Pacific Conference Series*, 150, 2-10.
- J. Rösch, "Lucien d’Azambuja (1884-1970)", 1970, *Solar Physics*, 15, 261-264.
- Vernazza, J.E., Avrett, E.H., Loeser, R., "Structure of the solar chromosphere. III. Models of the UEV brightness component of the quiet sun", 1981, *ApJS*, 45, 635.

11 - THE AUTHOR

Dr Jean-Marie Malherbe, born in 1956, is emeritus astronomer at Paris-Meudon observatory. He got the degrees of "*Docteur en astrophysique*" in 1983 and "*Docteur ès Sciences*" in 1987. He first worked on solar filaments and prominences using multi-wavelength observations. He used the spectrographs of the Meudon Solar Tower, the Pic du Midi Turret Dome, the German Vacuum Tower Telescope, THEMIS (Tenerife) and developed polarimeters. He proposed models and MHD 2D numerical simulations for prominence formation, including radiative cooling and magnetic reconnection. More recently, he worked on the quiet Sun, using satellites such as HINODE (JAXA), IRIS (NASA), and MHD 3D simulation results. He was responsible of the Meudon spectroheliograph from 1996 to 2023.

