

# **Air showers and their radio component**

Thierry Gousset (SubaTech, Nantes, France)

June 7, 2007

# Air showers

Refs:

— Allan, in: Progress in elementary particle and cosmic ray physics, p. 169 (North Holland, Amsterdam, 1971)

Refs:

- Allan, in: *Progress in elementary particle and cosmic ray physics*, p. 169 (North Holland, Amsterdam, 1971)
- Stanev, *High energy cosmic rays* (Springer, 2004)
- Gaisser, *Cosmic rays and particle physics* (Cambridge University Press, 1990)
- Nagano-Watson, *Rev Mod Phys* 72, 689 (2000)

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

≠ approaches

# Air showers and the primary radiation

## Length in m $\rightarrow$ length in g/cm<sup>2</sup>

Interaction of rays with matter described by various lengths  $\ell$   
 $\rightarrow$  mean free path for something to happen (coll, abs)  
probability of nothing to happen up to  $x$ :

$$\frac{dp}{p} = -\frac{dx}{\ell}$$

■  $1/\ell = \text{particle density} \times \text{cross section} = n \times \sigma$

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

$\neq$  approaches

## Length in m $\rightarrow$ length in g/cm<sup>2</sup>

Interaction of rays with matter described by various lengths  $\ell$   
 $\rightarrow$  mean free path for something to happen (coll, abs)  
probability of nothing to happen up to  $x$ :

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

$\neq$  approaches

$$\frac{dp}{p} = -\frac{dx}{\ell}$$

■  $1/\ell = \text{particle density} \times \text{cross section} = n \times \sigma$

■  $n \rightarrow n(x)$ , useful to use **depth of material  $X$**  such that  
 $dX = \rho(x)dx$

$$\frac{dp}{p} = -\frac{dX}{\lambda}$$

with  $\lambda$  in units of  $X$ , in practice g/cm<sup>2</sup>

85 g/cm<sup>2</sup> **vs** 1000 g/cm<sup>2</sup>

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

≠ approaches

- Earth's atmosphere = cosmic-ray shield
- High energy protons have interaction length in air
- Note:  $A$  for air (or 80% N+ 20% O)
- For a downward vertical path to sea level

$$\lambda_{pA} = 85 \text{ g/cm}^2$$

$$\int dX \approx 1000 \text{ g/cm}^2$$



# Cosmic rays and high energy physics

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

≠ approaches

$$p(E_p = 10^{17} \text{ eV}) + A \rightarrow X$$

$$s_{NN} = 2m_N c^2 E_p = O((10 \text{ TeV})^2), \text{ i.e. LHC}$$

# Cosmic rays and high energy physics

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

≠ approaches

$$p(E_p = 10^{17} \text{ eV}) + A \rightarrow X$$

$$s_{NN} = 2m_N c^2 E_p = O((10 \text{ TeV})^2), \text{ i.e. LHC}$$

■ Tevatron  $\rightarrow (2 \text{ TeV})^2$  and RHIC  $\rightarrow (200 \text{ GeV})^2$

$\rightarrow$  **need to extrapolate**

$\rightarrow$  hadronic models (more in Stanev sec 8.3)

# Secondary particles



## ■ $X$

- $\sim 10^2$  pions (20% something else) + target fragments + “original” baryon with a fraction of the initial energy
- pions are  $\pi^+$ ,  $\pi^-$  and  $\pi^0$

## ■ $\pi^0 \xrightarrow{99\%} 2\gamma$

- $c\tau_{\pi^0} = 25 \text{ nm}$ ;  $\pi^0$ 's desintegrate before reinteracting

## ■ $\gamma$ 's initiate the electromagnetic component of the shower

1. pair creation  $\gamma + A \rightarrow e^+e^- + X$
2. bremsstrahlung  $e + A \rightarrow e + \gamma + X$
3. repeat 1 and 2

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development

Shower extension

A first exercise

$\neq$  approaches

# Shower in space and time

- multiplicative process
- energy distributed among a vast number of secondary particles
- almost forward development

Air showers

Lengths

85 vs 1000

particle physics

secondary part

space and time

Time development


Shower extension

A first exercise

≠ approaches

time picture

shower  
history



Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

# Time development and energy distribution

# Time development

which  $N$ ?

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower


charge excess

Shower extension

A first exercise

$\neq$  approaches

- electrons and positrons
- charges
- charges above an energy threshold (in practice that of particle detection)



$N(X)$  trend results from competition

- **multiplicative processes**  $\Rightarrow dN > 0$  and  $E \searrow$
- **ionisation loss**  $\Rightarrow E \searrow$

# Heitler model

## Toy model for cascade development

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- $1 \rightarrow 2$  process, with daughter particles each carrying half the parent energy
- Branching at every step of length  $X_{1/2}$
- After the  $k$ th branching  $X = k \times X_{1/2}$ ,  $N = 2^k$  and the energy per particle is  $E_p/N$
- Assume branching process stops when  $E \leq E_C$

$$N_{\max} = \frac{E_p}{E_C}, \quad X_{\max} = X_{1/2} \log_2(E_p/E_C)$$

# $\gamma$ initiated shower

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

■  $\gamma + A \rightarrow e^+e^- + X$  and  $e + A \rightarrow e + \gamma + X$  are 1  $\rightarrow$  2 processes

■  $\approx$  same length scale 'radiation length' =  $X_0 \approx 40 \text{ g/cm}^2$

■  $X_{1/2} \approx \ln 2 \times X_0 = 30 \text{ g/cm}^2$

■ these branchings dominate for  $E > E_C$ , with a critical energy in air  $\approx 100 \text{ MeV}$

$$N_{\max} = \frac{E_\gamma}{100 \text{ MeV}}, \quad X_{\max} = 100 \text{ g/cm}^2 \times \log_{10}(E_\gamma/100 \text{ MeV})$$



# $\gamma$ initiated shower

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

■  $\gamma + A \rightarrow e^+e^- + X$  and  $e + A \rightarrow e + \gamma + X$  are 1  $\rightarrow$  2 processes

■  $\approx$  same length scale 'radiation length' =  $X_0 \approx 40 \text{ g/cm}^2$

■  $X_{1/2} \approx \ln 2 \times X_0 = 30 \text{ g/cm}^2$

■ these branchings dominate for  $E > E_C$ , with a critical energy in air  $\approx 100 \text{ MeV}$

$$N_{\max} = \frac{E_\gamma}{100 \text{ MeV}}, \quad X_{\max} = 100 \text{ g/cm}^2 \times \log_{10}(E_\gamma/100 \text{ MeV})$$

model misses energy loss by ionisation  $\Rightarrow N_{\max}$  overestimated

# Greisen parametrization

(Stanev p. 175)

$$N_e^\gamma = \frac{0.31}{\sqrt{\ln E_\gamma/E_C}} \exp \left[ \left(1 - \frac{3}{2} \ln s\right) X/X_0 \right], \quad s = \frac{3X}{X + 2X_{\max}}$$

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice


nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches



# $N_{\max}$ from total track length

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

■ Energy dissipated in ionisation loss; for relativistic particle the rate is  $dE/dX \approx -2 \text{ MeV/g/cm}^2$

■  $N(X)$  number of charged particles at depth  $X$

■ Energy dumped in  $[X, X + dX]$  slice

$$dE = (2 \text{ MeV/g/cm}^2) \times N(X)dX$$

# $N_{\max}$ from total track length

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- Energy dissipated in ionisation loss; for relativistic particle the rate is  $dE/dX \approx -2 \text{ MeV/g/cm}^2$

- $N(X)$  number of charged particles at depth  $X$

- Energy dumped in  $[X, X + dX]$  slice

$$dE = (2 \text{ MeV/g/cm}^2) \times N(X)dX$$

$\Rightarrow$

$$\int N(X)dX \approx \frac{E_p}{2 \text{ GeV}} \times 1000 \text{ g/cm}^2$$

# $N_{\max}$ from total track length

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- Energy dissipated in ionisation loss; for relativistic particle the rate is  $dE/dX \approx -2 \text{ MeV/g/cm}^2$

- $N(X)$  number of charged particles at depth  $X$

- Energy dumped in  $[X, X + dX]$  slice

$$dE = (2 \text{ MeV/g/cm}^2) \times N(X)dX$$

$\Rightarrow$

$$\int N(X)dX \approx \frac{E_p}{2 \text{ GeV}} \times 1000 \text{ g/cm}^2$$

( $\rightarrow$  fluorescence method, more in Nagano-Watson)

## $N_{\max}$ from total track length (cont'd)

$\int N(X)dX = N_{\max} \times \text{characteristic shower length}$

taking 1 atmospheric thickness:

$$N_{\max} = \frac{E_p}{2 \text{ GeV}}$$

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

# Energy spectrum (e-m component)

in Heitler model:

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- total track length associated with particles of energy greater than  $E = E_\gamma/2^k$   
 $= 30 \text{ g/cm}^2 \times 2^k (1/2 + 1/4 + \dots) \approx (E_\gamma/E) \times 30 \text{ g/cm}^2$

# Energy spectrum (e-m component)

in Heitler model:

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- total track length associated with particles of energy greater than  $E = E_\gamma/2^k$   
 $= 30 \text{ g/cm}^2 \times 2^k (1/2 + 1/4 + \dots) \approx (E_\gamma/E) \times 30 \text{ g/cm}^2$
- an electron with  $E = E_C$  loses it in one radiation length
- total track length associated with particles of energy lower than  $E_C = (E_\gamma/E_C) \times 40 \text{ g/cm}^2$



# Energy spectrum (e-m component)

in Heitler model:

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- total track length associated with particles of energy greater than  $E = E_\gamma/2^k$   
 $= 30 \text{ g/cm}^2 \times 2^k (1/2 + 1/4 + \dots) \approx (E_\gamma/E) \times 30 \text{ g/cm}^2$
- an electron with  $E = E_C$  loses it in one radiation length
- total track length associated with particles of energy lower than  $E_C = (E_\gamma/E_C) \times 40 \text{ g/cm}^2$
- more weight to low energy in actual fact

$$\int_{>E} N dX \approx 40 \text{ g/cm}^2 \times \frac{E_\gamma}{E_C} \times \frac{30 \text{ MeV}}{E + 30 \text{ MeV}}$$

# Energy spectrum (e-m component)

in Heitler model:

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- total track length associated with particles of energy greater than  $E = E_\gamma/2^k$   
 $= 30 \text{ g/cm}^2 \times 2^k (1/2 + 1/4 + \dots) \approx (E_\gamma/E) \times 30 \text{ g/cm}^2$
- an electron with  $E = E_C$  loses it in one radiation length
- total track length associated with particles of energy lower than  $E_C = (E_\gamma/E_C) \times 40 \text{ g/cm}^2$
- more weight to low energy in actual fact

$$\int_{>E} N dX \approx 40 \text{ g/cm}^2 \times \frac{E_\gamma}{E_C} \times \frac{30 \text{ MeV}}{E + 30 \text{ MeV}}$$

this is for the whole shower  $\rightarrow$  at and around maximum

$$N(> E)/N$$

# Hadronic component

■  $\pi^\pm$ :  $c\tau = 8 \text{ m}$

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

—  $\lambda_{\pi A} = 120 \text{ g/cm}^2 \rightarrow \ell_{\pi A} \approx 1 \text{ km}$  for  
 $n(z=0) = 1 \text{ mg/cm}^3$

— at high energy pions **reinteract**

— otherwise they **decay**  $\rightarrow \mu\nu$ ; muons (only lose  
 $2 \text{ MeV/g/cm}^2$ )  $\rightarrow$  direct information on pions

# Hadronic component

■  $\pi^\pm$ :  $c\tau = 8 \text{ m}$

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

—  $\lambda_{\pi A} = 120 \text{ g/cm}^2 \rightarrow \ell_{\pi A} \approx 1 \text{ km}$  for  
 $n(z=0) = 1 \text{ mg/cm}^3$

— at high energy pions **reinteract**

— otherwise they **decay**  $\rightarrow \mu\nu$ ; muons (only lose  
 $2 \text{ MeV/g/cm}^2$ )  $\rightarrow$  direct information on pions

■  $\pi^0$ : estimate of  $X_{\max}$  and  $N_{\max}$  for proton induced shower assuming that the e.m. showers are initiated by 1st generation  $\pi^0$ 's

$$X_{\max} = \lambda_{pA} + X_0 \ln \left[ \frac{(1-K) E_p}{2 \langle m \rangle E_C} \right], \quad N_{\max} = \frac{(1-K) E_p}{3 E_C}$$

## in practice

### average behavior (adjusted with Monte-Carlo)

Gaisser-Hillas formula (Stanev p. 186)

$$N(X) = N_{\max} \left( \frac{X - X_1}{X_{\max} - \lambda} \right)^{X_{\max}/\lambda - 1} \exp - \left( \frac{X - X_1}{\lambda} \right)$$

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

## in practice

### average behavior (adjusted with Monte-Carlo)

Gaisser-Hillas formula (Stanev p. 186)

$$N(X) = N_{\max} \left( \frac{X - X_1}{X_{\max} - \lambda} \right)^{X_{\max}/\lambda - 1} \exp - \left( \frac{X - X_1}{\lambda} \right)$$

+ fluctuations:

- on  $X_1 \rightarrow X_{\max}$
- on shape and  $N_{\max}$ : individual realizations of first hadronic collisions (inelasticity, multiplicity, energy of secondaries)

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

# Nucleus vs proton initiated shower

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

charge excess

Shower extension

A first exercise

$\neq$  approaches

- in the superposition approximation: nucleus =  $A \times$  independent nucleons with energy  $E_p/A$
- nucleus shower =  $A \times$  nucleon showers
- shift of  $X_{\max}$ :  $X_{\max}(E_p, A) = X_{\max}(E_p/A, p)$
- less shower to shower fluctuation

# Negative charge excess

- below  $E_C$



**delta rays** (Compton recoil)

- positron annihilate in flight

→ 10–20%  $e^-$  excess in the energy range below  $E_C$

Air showers

Time development

$N(X)$

Heitler model

$\gamma$  shower

Greisen

$N_{\max}$

energy spectrum

hadron content

in practice

nucleus shower

**charge excess**

Shower extension

A first exercise

≠ approaches



Air showers

Time development

Shower extension

Multiple scattering  
NKG

A first exercise

≠ approaches

# Lateral spread and longitudinal dispersion

# Multiple scattering

- **Emission** → spread of hadrons

- **Multiple scattering** → spread of electrons

spread of hadrons limited to a few meters

Air showers

Time development

Shower extension

Multiple scattering

NKG

A first exercise

≠ approaches

# Multiple scattering

- **Emission** → spread of hadrons

- **Multiple scattering** → spread of electrons

spread of hadrons limited to a few meters

## electrons

- typical scattering angle  $\theta \sim 1/\gamma$

- $\theta^2(n) = n \times (1/\gamma)^2$

- proportion of scatterings with radiation  $\sim 1/\alpha$

$$\Rightarrow d\theta^2 = (E_s/E)^2 dX/X_0, \text{ with } E_s = 4\pi m_e c^2 / \alpha = 21 \text{ MeV}$$

Air showers

Time development

Shower extension

Multiple scattering

NKG

A first exercise

≠ approaches

## Multiple scattering (cont'd)

including energy loss  $X_i \rightarrow X_f$ ,  $E(X) = E(X_f) \times e^{\frac{X_f - X}{X_0}}$

Air showers

Time development

Shower extension

Multiple scattering

NKG

A first exercise

≠ approaches

$$X' = X_f - X,$$

$$\theta^2(i \rightarrow f) = \int d\theta^2 = \frac{E_s^2}{E_f^2} \int_0^{X_i - X_f} e^{-2X'/X_0} \frac{dX'}{X_0}, \quad (E_f > E_C)$$

## Multiple scattering (cont'd)

including energy loss  $X_i \rightarrow X_f$ ,  $E(X) = E(X_f) \times e^{\frac{X_f - X}{X_0}}$

Air showers

Time development

Shower extension

Multiple scattering

NKG

A first exercise

≠ approaches

$$X' = X_f - X,$$

$$\theta^2(i \rightarrow f) = \int d\theta^2 = \frac{E_s^2}{E_f^2} \int_0^{X_i - X_f} e^{-2X'/X_0} \frac{dX'}{X_0}, \quad (E_f > E_C)$$

and lateral displacement

$$D^2(i \rightarrow f) = \int X'^2 d\theta^2 \Rightarrow D = \frac{10 \text{ MeV}}{E} X_0 \quad (E > E_C)$$

i.e., 40 m at 100 MeV at sea level

## Multiple scattering (cont'd)

including energy loss  $X_i \rightarrow X_f$ ,  $E(X) = E(X_f) \times e^{\frac{X_f - X}{X_0}}$

Air showers

Time development

Shower extension

Multiple scattering

NKG

A first exercise

≠ approaches

$$X' = X_f - X,$$

$$\theta^2(i \rightarrow f) = \int d\theta^2 = \frac{E_s^2}{E_f^2} \int_0^{X_i - X_f} e^{-2X'/X_0} \frac{dX'}{X_0}, \quad (E_f > E_C)$$

and lateral displacement

$$D^2(i \rightarrow f) = \int X'^2 d\theta^2 \Rightarrow D = \frac{10 \text{ MeV}}{E} X_0 \quad (E > E_C)$$

i.e., **40 m at 100 MeV at sea level**

→ multiple scattering longitudinal lag **3 m**, also ↘ with ↗ energy

## Lateral distribution

flux of electrons given by NKG formula (Gaisser p 226, Stanev p 179)

Air showers

Time development

Shower extension

Multiple scattering

**NKG**

$$n_e(r, X) = N_e(X) \frac{C}{r r_1} \left( \frac{r}{r_1} \right)^{s-1} \left( 1 + \frac{r}{r_1} \right)^{s-9/2},$$

with

$$r_1 = \frac{E_s}{E_C} \frac{X_0}{\rho_{\text{air}}}$$

A first exercise

≠ approaches

# From showers to electric fields



Air showers

Time development

Shower extension

A first exercise

Detour

Coherence

≠ approaches

# A first exercise

## A detour: Cerenkov light in air

consider Cerenkov radiation of a charge particle ( $q = Ze$ )  
energy spectrum per unit length

Air showers

Time development

Shower extension

A first exercise

**Detour**

Coherence

≠ approaches

$$\frac{d^2 E_C}{dL d\omega} = \alpha Z^2 \sin^2 \theta_C \frac{\omega}{\hbar c}$$

## A detour: Cerenkov light in air

consider Cerenkov radiation of a charge particle ( $q = Ze$ )  
energy spectrum per unit length

Air showers

Time development

Shower extension

A first exercise

**Detour**

Coherence

≠ approaches

$$\frac{d^2 E_C}{dL d\omega} = \alpha Z^2 \sin^2 \theta_C \frac{\omega}{\hbar c}$$

- Cerenkov in air  $\theta_C \ll 1$
- vertical downward moving particle
- trajectory bit of length  $\Delta z$  around  $z_0$  shines on a ring of mean radius  $z_0 \theta_C$  and width  $\Delta z \theta_C$

$$\frac{\Delta E_C}{2\pi z_0 \Delta z \theta_C^2} = \alpha Z^2 \frac{\omega \Delta \omega}{\hbar c z_0}$$

# Cerenkov radio

$$\Rightarrow dE_C/dS \sim 10^2 \text{ MeV/m}^2 \text{ using } hc = 1.24 \text{ eV } \mu\text{m}, Z = 1, \\ z_0 = 4 \text{ km}, N_e = 5 \cdot 10^7, \lambda = 0.6 \mu\text{m} \text{ and } \Delta\lambda = 0.4 \mu\text{m}$$

Air showers

Time development

Shower extension

A first exercise

Detour

Coherence

≠ approaches

# Cerenkov radio

$$\Rightarrow dE_C/dS \sim 10^2 \text{ MeV/m}^2 \text{ using } hc = 1.24 \text{ eV } \mu\text{m}, Z = 1, \\ z_0 = 4 \text{ km}, N_e = 5 \cdot 10^7, \lambda = 0.6 \mu\text{m} \text{ and } \Delta\lambda = 0.4 \mu\text{m}$$

Air showers

Time development

Shower extension

A first exercise

Detour

Coherence

≠ approaches

radio (decametric)

■ divide  $\omega$  by  $\sim 10^7$  and  $\Delta\omega$  by  $\sim 10^7$

■ take  $A_e \sim 10 \text{ m}^2$

$$\Rightarrow \Delta E_C = 10^{-5} \text{ eV}$$

# Cerenkov radio

$$\Rightarrow dE_C/dS \sim 10^2 \text{ MeV/m}^2 \text{ using } hc = 1.24 \text{ eV } \mu\text{m}, Z = 1, \\ z_0 = 4 \text{ km}, N_e = 5 \cdot 10^7, \lambda = 0.6 \mu\text{m} \text{ and } \Delta\lambda = 0.4 \mu\text{m}$$

Air showers

Time development

Shower extension

A first exercise

Detour

Coherence

≠ approaches

radio (decametric)

■ divide  $\omega$  by  $\sim 10^7$  and  $\Delta\omega$  by  $\sim 10^7$

■ take  $A_e \sim 10 \text{ m}^2$

$$\Rightarrow \Delta E_C = 10^{-5} \text{ eV}$$

■ much too small since galactic noise gives

$$k_B T \times \Delta\nu \times \Delta t \rightarrow 2.5 \text{ eV} \times 40 \text{ MHz} \times 10 \text{ ns} = 1 \text{ eV}$$

# Coherence: a must in radio

**solution: replace  $N_e \rightarrow N_e^2$ ; incoherent  $\rightarrow$  coherent**

Air showers

Time development

Shower extension

A first exercise

Detour

Coherence

$\neq$  approaches

■ at first:  $N_- = N_+ \Rightarrow$  no field at all

■ but *systematic* charge separation by

1. **earth magnetic field** (and  $E$  field in thunderstorms)
2. **elementary processes below  $E_C$**   $\rightarrow$  negative charge excess

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

## Various approaches



# Overview

Air showers

Time development

Shower extension

A first exercise

≠ approaches

**Overview**

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

- $\sum_{k=1}^N \vec{E}(t, A)$  with  $\vec{E}$  single-charge electric field taken from textbook → Monte-Carlo based approach
- $\vec{E}[\rho, \vec{j}]$
- Feynman formula for relativistic charges:

$$\vec{E} = \frac{-q}{4\pi\epsilon_0 c^2} \vec{e}_{r'}$$

# From individual charges

More thorough study to date: T Huege, H Falcke, *Astronomy & Astrophysics* 412, 19 (2003); *Astronomy & Astrophysics* 430, 779 (2005); *Astropart. Phys.* 24, 116 (2005); T Huege et al, *Astropart. Phys.* 27, 392 (2007)

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

## From individual charges

More thorough study to date: T Huege, H Falcke, *Astronomy & Astrophysics* 412, 19 (2003); *Astronomy & Astrophysics* 430, 779 (2005); *Astropart. Phys.* 24, 116 (2005); T Huege et al, *Astropart. Phys.* 27, 392 (2007)

Building block:

$$\vec{E}_q(t, A) = \frac{q}{4\pi\epsilon c^2} \frac{\vec{R} \wedge [(\vec{R} - R\vec{v}/c) \wedge \vec{a}]}{||R - \vec{R} \cdot \vec{v}/c||^3}$$

with

$$\vec{a} = \frac{q\vec{v} \wedge \vec{B}}{\gamma m_e}$$

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

# From charges and currents

show electromagnetic field as a standard electromagnetism exercise

$$(\vec{E}, \vec{B}) = F[\rho, \vec{j}]$$

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

# From charges and currents

show electromagnetic field as a standard electromagnetism exercise

$$(\vec{E}, \vec{B}) = F[\rho, \vec{j}]$$

how to carry out such a program ?

## ■ Kahn and Lerche approach

- ringlike geometry
- no shower evolution (contribution around  $N_{\max}$ ) + estimate for shower decay
- formulation in Fourier space
- geomagnetic contribution  $>$  Askaryan effect (charge excess)

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

# From charges and currents

show electromagnetic field as a standard electromagnetism exercise

$$(\vec{E}, \vec{B}) = F[\rho, \vec{j}]$$

how to carry out such a program ?

## ■ Kahn and Lerche approach

- ringlike geometry
- no shower evolution (contribution around  $N_{\max}$ ) + estimate for shower decay
- formulation in Fourier space
- geomagnetic contribution  $>$  Askaryan effect (charge excess)

- more realistic model, numerical implementation... not (yet) followed

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

# Time scales

Air showers

Time development

Shower extension

A first exercise

≠ approaches


Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$



■ particle  $Q$  moves at  $\approx c$

■  $Q = B$  at  $t = 0$

■  $ct_i = \sqrt{d_{iB}^2 + b^2} - d_{iB}$


■  $\Delta t_{12} \approx 0.4 b^2$  (small  $\theta$ )

■  $\Delta t_{2B} = 3.3 b$

(time in  $\mu\text{s}$  and distance in km)

# Time scales

- Air showers
- Time development
- Shower extension
- A first exercise
- ≠ approaches
- Overview
- $\Sigma \vec{E}$
- $\vec{E}[\rho, \vec{j}]$
- Time scales**
- Large  $b$



- particle  $Q$  moves at  $\approx c$

- $Q = B$  at  $t = 0$

- $ct_i = \sqrt{d_{iB}^2 + b^2} - d_{iB}$

- $\Delta t_{12} \approx 0.4 b^2$  (small  $\theta$ )

- $\Delta t_{2B} = 3.3 b$

(time in  $\mu s$  and distance in km)


- Doppler distortion: **fast rise and slow decay**

- $v \approx c$  valid at  $\theta \gg |c - v|$



# Time scales

- Air showers
- Time development
- Shower extension
- A first exercise
- ≠ approaches
- Overview
- $\Sigma \vec{E}$
- $\vec{E}[\rho, \vec{j}]$
- Time scales**
- Large  $b$



- particle  $Q$  moves at  $\approx c$

- $Q = B$  at  $t = 0$

- $ct_i = \sqrt{d_{iB}^2 + b^2} - d_{iB}$

- $\Delta t_{12} \approx 0.4 b^2$  (small  $\theta$ )

- $\Delta t_{2B} = 3.3 b$

(time in  $\mu\text{s}$  and distance in km)

- Doppler distortion: **fast rise and slow decay**

- $v \approx c$  valid at  $\theta \gg |c - v|$

- $\Delta t_{32} \approx l/c + br/(cd_{2B})$ , both terms  $< 30$  ns

# Large impact parameters

⇒ at large  $b \rightarrow$  (distorted) image of  $N(X)$

- pointlike approximation: all timescales but obliquity set to 0

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

# Large impact parameters

⇒ at large  $b \rightarrow$  (distorted) image of  $N(X)$

Air showers

Time development

Shower extension

A first exercise

≠ approaches

Overview


$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales

Large  $b$

- pointlike approximation: all timescales but obliquity set to 0



$$-ct' \gg b^2/2 \gg ct$$

$$ct ct' \approx -b^2/2,$$

$$E(t, A) = \frac{e N_{ee}(t') a_T}{4\pi\epsilon c^2} \frac{b^2}{2 (ct)^3}$$

$$a_T = \frac{e c B \sin \alpha}{\gamma m_e}$$

# Large impact parameters (cont'd)

(vertical,  $10^{19}$  eV, 700 m)

Air showers

Time development

Shower extension

A first exercise

≠ approaches



Overview

$\Sigma \vec{E}$

$\vec{E}[\rho, \vec{j}]$

Time scales



Large  $b$



# Large impact parameters (cont'd)

(vertical,  $10^{19}$  eV, 700 m)

- Air showers
- Time development
- Shower extension
- A first exercise
- $\neq$  approaches
- Overview
- $\Sigma \vec{E}$
- $\vec{E}[\rho, \vec{j}]$
- Time scales
- Large  $b$



makes it possible

- problem inversion
- discussion of antenna spacing for a giant array